

THE GROUP OF UNITAL C^* -EXTENSIONS

VLADIMIR MANUILOV AND KLAUS THOMSEN

ABSTRACT. Let A and B be separable C^* -algebras, A unital and B stable. It is shown that there is a natural six-terms exact sequence which relates the group which arises by considering *all* semi-split extensions of A by B to the group which arises by restricting the attention to *unital* semi-split extensions of A by B . The six-terms exact sequence is an unpublished result of G. Skandalis.

Let A, B be separable C^* -algebras, B stable. As is well-known the C^* -algebra extensions of A by B can be identified with $\text{Hom}(A, Q(B))$, the set of $*$ -homomorphisms $A \rightarrow Q(B)$ where $Q(B) = M(B)/B$ is the generalized Calkin algebra. Two extensions $\varphi, \psi : A \rightarrow Q(B)$ are *unitarily equivalent* when there is a unitary $u \in M(B)$ such that $\text{Ad } q(u) \circ \psi = \varphi$, where $q : M(B) \rightarrow Q(B)$ is the quotient map. The unitary equivalence classes of extensions of A by B have an abelian semi-group structure thanks to the stability of B : Choose isometries $V_1, V_2 \in M(B)$ such that $V_1 V_1^* + V_2 V_2^* = 1$, and define the sum $\varphi \oplus \psi : A \rightarrow Q(B)$ of $\varphi, \psi \in \text{Hom}(A, Q(B))$ by

$$(\psi \oplus \varphi)(a) = \text{Ad } q(V_1) \circ \psi(a) + \text{Ad } q(V_2) \circ \varphi(a). \quad (1)$$

The isometries, V_1 and V_2 , are fixed in the following. An extension $\varphi : A \rightarrow Q(B)$ is *split* when there is a $*$ -homomorphism $\pi : A \rightarrow M(B)$ such that $\varphi = q \circ \pi$. To trivialize the split extensions we declare two extensions $\varphi, \psi : A \rightarrow Q(B)$ to be *stably equivalent* when there is a split extension π such that $\psi \oplus \pi$ and $\varphi \oplus \pi$ are unitarily equivalent. This is an equivalence relation because the sum (1) of two split extensions is itself split. We denote by $\text{Ext}(A, B)$ the semigroup of stable equivalence classes of extensions of A by B . It was proved in [Th], as a generalization of results of Kasparov, that there exists an *absorbing split extension* $\pi_0 : A \rightarrow Q(B)$, i.e. a split extension with the property that $\pi_0 \oplus \pi$ is unitarily equivalent to π_0 for every split extension π . Thus two extensions φ, ψ are stably equivalent if and only if $\varphi \oplus \pi_0$ and $\psi \oplus \pi_0$ are unitarily equivalent. The classes of stably equivalent extensions of A by B is an abelian semigroup $\text{Ext}(A, B)$ in which any split extension (like 0) represents the neutral element. As is well-documented the semi-group is generally not a group, and we denote by

$$\text{Ext}^{-1}(A, B)$$

the abelian group of invertible elements in $\text{Ext}(A, B)$. It is also well-known that this group is one way of describing the KK -groups of Kasparov. Specifically, $\text{Ext}^{-1}(A, B) = KK(SA, B) = KK(A, SB)$.

Assume now that A is unital. It is then possible, and sometimes even advantageous, to restrict attention to unital extensions of A by B , i.e. to short exact sequences

$$0 \longrightarrow B \longrightarrow E \longrightarrow A \longrightarrow 0 \quad (2)$$

of C^* -algebras with E is unital, or equivalently to $*$ -homomorphisms $A \rightarrow Q(B)$ that are unital. The preceding definitions are all amenable to such a restriction, if done consistently. Specifically, we say that a unital extension $\varphi : A \rightarrow Q(B)$ is *unittally split* when there is a unital $*$ -homomorphism $\pi : A \rightarrow M(B)$ such that $\varphi = q \circ \pi$. The sum (1) of two unital extensions is again unital, and we say that two unital extensions $\varphi, \psi : A \rightarrow Q(B)$ are *unittally stably equivalent* when there is a unital split extension π such that $\psi \oplus \pi$ and $\varphi \oplus \pi$ are unitarily equivalent. It was proved in [Th] that there always exists a *unittally absorbing split extension* $\pi_0 : A \rightarrow Q(B)$, i.e. a unittally split extension with the property that $\pi_0 \oplus \pi$ is unitarily equivalent to π_0 for every unittally split extension π . Thus two unital extensions φ, ψ are unittally stably equivalent if and only if $\varphi \oplus \pi_0$ and $\psi \oplus \pi_0$ are unitarily equivalent. The classes of unittally stably equivalent extensions of A by B is an abelian semi-group which we denote by $\text{Ext}_{\text{unital}}(A, B)$. The unittally absorbing split extension π_0 , or any other unittally split extension, represents the neutral element of $\text{Ext}_{\text{unital}}(A, B)$, and we denote by

$$\text{Ext}_{\text{unital}}^{-1}(A, B)$$

the abelian group of invertible elements in $\text{Ext}_{\text{unital}}(A, B)$. As we shall see there is a difference between $\text{Ext}_{\text{unital}}^{-1}(A, B)$ and $\text{Ext}^{-1}(A, B)$ arising from the fact that while the class in $\text{Ext}^{-1}(A, B)$ of a unital extension $A \rightarrow Q(B)$ can not be changed by conjugating it with a unitary from $Q(B)$, its class in $\text{Ext}_{\text{unital}}^{-1}(A, B)$ can. In a sense the main result of this note is that this is the only way in which the two groups differ.

Note that there is a group homomorphism

$$\text{Ext}_{\text{unital}}^{-1}(A, B) \rightarrow \text{Ext}^{-1}(A, B),$$

obtained by forgetting the word 'unital'. It will be shown that this forgetful map fits into a six-terms exact sequence

$$\begin{array}{ccccc} K_0(B) & \xrightarrow{u_0} & \text{Ext}_{\text{unital}}^{-1}(A, B) & \longrightarrow & \text{Ext}^{-1}(A, B) \\ \uparrow i_0^* & & & & \downarrow i_1^* \\ \text{Ext}^{-1}(A, SB) & \longleftarrow & \text{Ext}_{\text{unital}}^{-1}(A, SB) & \xleftarrow{u_1} & K_1(B) \end{array} \quad (3)$$

where SB is the suspension of B , i.e. $SB = C_0(0, 1) \otimes B$, and the maps u_k and i_k^* , $k = 0, 1$, will be defined shortly. This six-terms exact sequence is mentioned in 10.11 of [S], but the proof was never published.

Fix a unittally absorbing $*$ -homomorphism $\alpha_0 : A \rightarrow M(B)$, which exists by Theorem 2.4 of [Th]. It follows then from Theorem 2.1 of [Th] that $\alpha = q \circ \alpha_0$ is a unittally absorbing split extension as defined above.

Lemma 1. *The $*$ -homomorphisms $\text{Ad } V_1 \circ \alpha_0 : A \rightarrow M(B)$ and $(\begin{smallmatrix} \alpha_0 & 0 \\ 0 & 0 \end{smallmatrix}) : A \rightarrow M(M_2(B))$ are both absorbing.*

Proof. There is a $*$ -isomorphism $M(M_2(B)) = M_2(M(B)) \rightarrow M(B)$ given by

$$\begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix} \mapsto V_1 m_{11} V_1^* + V_1 m_{12} V_2^* + V_2 m_{21} V_1^* + V_2 m_{22} V_2^*,$$

which sends $M_2(B)$ to B and $(\begin{smallmatrix} \alpha_0 & 0 \\ 0 & 0 \end{smallmatrix})$ to $\text{Ad } V_1 \circ \alpha_0$, so it suffices to show that the latter is an absorbing $*$ -homomorphism. By definition, cf. Definition 2.6 of [Th], we must show that the unital $*$ -homomorphism $A \oplus \mathbb{C} \ni (a, \lambda) \mapsto V_1 \alpha_0(a) V_1^* + \lambda V_2 V_2^*$ is unittally absorbing.

For this we check that it has property 1) of Theorem 2.1 of [Th]. So let $\varphi : A \oplus \mathbb{C} \rightarrow B$ be a completely positive contraction. Since α_0 has property 1), there is a sequence $\{W_n\}$ in $M(B)$ such that $\lim_{n \rightarrow \infty} W_n^* b = 0$ for all $b \in B$ and $\lim_{n \rightarrow \infty} W_n^* \alpha_0(a) W_n = \varphi(a)$ for all $a \in A$. Since B is stable there is a sequence $\{S_n\}$ of isometries in $M(B)$ such that $\lim_{n \rightarrow \infty} S_n^* b = 0$ for all $b \in B$. Set

$$T_n = V_1 W_n + V_2 S_n \varphi(0, 1)^{\frac{1}{2}}.$$

Then $\lim_{n \rightarrow \infty} T_n b = 0$ for all $b \in B$, and

$$T_n^* (V_1 \alpha_0(a) V_1^* + \lambda V_2 V_2^*) T_n = W_n^* \alpha_0(a) W_n + \varphi(0, \lambda)$$

for all n . Since the last expression converges to $\varphi(a, \lambda)$ as n tends to infinity, the proof is complete. $\square$

Set

$$C_\alpha = \{m \in M_2(M(B)) : m \begin{pmatrix} \alpha_0(a) & 0 \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} \alpha_0(a) & 0 \\ 0 & 0 \end{pmatrix} m \in M_2(B) \ \forall a \in A\}$$

and

$$A_\alpha = \{m \in C_\alpha : m \begin{pmatrix} \alpha_0(a) & 0 \\ 0 & 0 \end{pmatrix} \in M_2(B) \ \forall a \in A\}.$$

We can define a $*$ -homomorphism $C_\alpha \rightarrow \alpha(A)' \cap Q(B)$ such that

$$\begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix} \mapsto q(m_{11}).$$

Then kernel is then A_α , so we have a $*$ -isomorphism $C_\alpha/A_\alpha \simeq \alpha(A)' \cap Q(B)$. By Lemma 1 (α_0) is an absorbing $*$ -homomorphism, so we conclude from Theorem 3.2 of [Th] that there is an isomorphism

$$K_1(\alpha(A)' \cap Q(B)) \simeq KK(A, B). \quad (4)$$

Since the unital $*$ -homomorphism $\mathbb{C} \rightarrow M(B)$ is unitaly absorbing, this gives us also the well-known isomorphism

$$K_1(Q(B)) \simeq KK(\mathbb{C}, B). \quad (5)$$

Let $i : \mathbb{C} \rightarrow A$ be the unital $*$ -homomorphism. For convenience we denote the map $K_1(\alpha(A)' \cap Q(B)) \rightarrow K_1(Q(B))$ induced by the inclusion $\alpha(A)' \cap Q(B) \subseteq Q(B)$ by i^* . It is then easy to check that the isomorphisms (4) and (5) match up to make the diagram

$$\begin{array}{ccc} K_1(\alpha(A)' \cap Q(B)) & \xrightarrow{i^*} & K_1(Q(B)) \\ \downarrow & & \downarrow \\ KK(A, B) & \xrightarrow{i^*} & KK(\mathbb{C}, B) \end{array} \quad (6)$$

commute.

Let v be a unitary in $M_n(Q(B))$. By composing the $*$ -homomorphism $\text{Ad } v \circ (1_n \otimes \alpha) : A \rightarrow M_n(Q(B))$ with an isomorphism $M_n(Q(B)) \simeq Q(B)$ which is canonical in the sense that it arises from an isomorphism $M_n(B) \simeq B$, we obtain a unital extension $e(v) : A \rightarrow Q(B)$ of A by B . By use of a unitary lift of $(v \ v^*)$ one sees that $e(v) \oplus e(v^*)$ is split, proving that $e(v)$ represents an element in $\text{Ext}_{\text{unital}}^{-1}(A, B)$. If $v_t, t \in [0, 1]$, is a norm-continuous path of unitaries in $M_n(Q(B))$ there is a partition $0 = t_0 < t_1 < t_2 < \dots < t_N = 1$ of $[0, 1]$ such that $v_{t_i} v_{t_{i+1}}^*$ is in the connected component of 1 in the unitary group of $M_n(Q(B))$

and hence has a unitary lift to $M_n(M(B))$. It follows that $e(v_0) = e(v_1)$, and it is then clear that the construction gives us a group homomorphism

$$u : K_1(Q(B)) \rightarrow \text{Ext}_{\text{unital}}^{-1}(A, B).$$

Lemma 2. *The sequence*

$$\begin{array}{ccccc} K_1(Q(B)) & \xrightarrow{u} & \text{Ext}_{\text{unital}}^{-1}(A, B) & \longrightarrow & \text{Ext}^{-1}(A, B) \\ & \uparrow i^* & & & \downarrow i^* \\ K_1(\alpha(A)' \cap Q(B)) & & & & \text{Ext}^{-1}(\mathbb{C}, B) \end{array} \quad (7)$$

is exact.

Proof. Exactness at $K_1(Q(B))$: If v is a unitary in $M_n(\alpha(A)' \cap Q(B))$, the extension $\text{Ad } v \circ (1_n \otimes \alpha) = 1_n \otimes \alpha$ (of A by $M_n(B)$) is split, proving that $u \circ i^* = 0$. To show that $\ker u \subseteq \text{im } i^*$, let $v \in Q(B)$ be a unitary such that $u[v] = 0$. Then $\text{Ad } v \circ \alpha \oplus \alpha$ is unitarily equivalent to $\alpha \oplus \alpha$, which means that there is a unitary $S \in M(M_2(B))$ such that

$$\text{Ad} \left((\text{id}_{M_2(\mathbb{C})} \otimes q)(S) \begin{pmatrix} v & \\ & 1 \end{pmatrix} \right) \begin{pmatrix} \alpha(a) & \\ & \alpha(a) \end{pmatrix} = \begin{pmatrix} \alpha(a) & \\ & \alpha(a) \end{pmatrix} \quad (8)$$

for all $a \in A$. Since the unitary group of $M(M_2(B))$ is normconnected by [M] or [CH], the unitary $\begin{pmatrix} v & \\ & 1 \end{pmatrix}$ is homotopic to $(\text{id}_{M_2(\mathbb{C})} \otimes q)(S) \begin{pmatrix} v & \\ & 1 \end{pmatrix}$ which is in $M_2(\alpha(A)' \cap Q(B))$ by (8). This implies that $[v] \in \text{im } i^*$. The same argument works when v is a unitary $M_n(Q(B))$ for some $n \geq 2$.

Exactness at $\text{Ext}_{\text{unital}}^{-1}(A, B)$: For any unitary $v \in Q(B)$,

$$(\text{Ad } v \circ \alpha) \oplus 0 = \text{Ad} \left((\text{id}_{M_2(\mathbb{C})} \otimes q)(T) \right) \circ (\alpha \oplus 0),$$

where $T \in M_2(M(B))$ is a unitary lift of $\begin{pmatrix} v & \\ & v^* \end{pmatrix}$. Hence $[\text{Ad } v \circ \alpha] = 0$ in $\text{Ext}^{-1}(A, B)$. The same argument works when v is a unitary $M_n(Q(B))$ for some $n \geq 2$, and we conclude that the composition $K_1(Q(B)) \rightarrow \text{Ext}_{\text{unital}}^{-1}(A, B) \rightarrow \text{Ext}^{-1}(A, B)$ is zero. Let $\varphi : A \rightarrow Q(B)$ be a unital extension such that $[\varphi] = 0$ in $\text{Ext}^{-1}(A, B)$. By Lemma 1 this means that there is a unitary $T \in M(M_3(B))$ such that

$$\text{Ad} \left((\text{id}_{M_3(\mathbb{C})} \otimes q)(T) \right) \circ \begin{pmatrix} \varphi & & \\ & \alpha & \\ & & 0 \end{pmatrix} = \begin{pmatrix} \alpha & & \\ & \alpha & \\ & & 0 \end{pmatrix}.$$

It follows that $(\text{id}_{M_3(\mathbb{C})} \otimes q)(T) = \begin{pmatrix} V & & \\ & r & \end{pmatrix}$ for some unitaries $V \in M_2(Q(B))$ and $r \in Q(B)$. Hence

$$\begin{pmatrix} \varphi & \alpha \end{pmatrix} = \text{Ad } V^* \circ \begin{pmatrix} \alpha & \alpha \end{pmatrix}.$$

Thus $[\varphi] = u[V^*]$.

Exactness at $\text{Ext}^{-1}(A, B)$: It is obvious that i^* kills the image of $\text{Ext}_{\text{unital}}^{-1}(A, B)$, so consider an invertible extension $\varphi : A \rightarrow Q(B)$ such that $[\varphi \circ i] = 0$ in $\text{Ext}^{-1}(\mathbb{C}, B)$. By Lemma 1, applied with $A = \mathbb{C}$, this means that there is a unitary $T \in M_3(M(B))$ such that

$$(\text{id}_{M_3(\mathbb{C})} \otimes q)(T) \begin{pmatrix} \varphi(1) & & \\ & 1 & \\ & & 0 \end{pmatrix} (\text{id}_{M_3(\mathbb{C})} \otimes q)(T^*) = \begin{pmatrix} 0 & & \\ & 1 & \\ & & 0 \end{pmatrix}. \quad (9)$$

Set $\psi = \varphi \oplus \alpha \oplus 0$. It follows from (9) that there are isometries $W_1, W_2, W_3 \in M(B)$ and a unitary $u \in M(B)$ such that $W_i^* W_j = 0, i \neq j$, $W_1 W_1^* + W_2 W_2^* + W_3 W_3^* = 1$ and $\text{Ad } q(u) \circ \psi(1) = q(W_2 W_2^*)$. Then $\text{Ad } q(u) \circ \psi + \text{Ad } q(W_1) \circ \alpha + \text{Ad } q(W_3) \circ \alpha$ is a unital extension which is invertible because it admits a completely positive contractive lifting

to $M(B)$ since ψ does, cf. [A]. As it represents the same class in $\text{Ext}^{-1}(A, B)$ as φ , the proof is complete. $\square$

In order to complete the sequence of Lemma 2, let $i_1^* : \text{Ext}^{-1}(A, B) \rightarrow K_1(B)$ be the composition

$$\text{Ext}^{-1}(A, B) \longrightarrow K_1(\alpha(A)' \cap Q(B)) \xrightarrow{i^*} K_1(Q(SB)) \longrightarrow K_1(B), \quad (10)$$

where the first map is the isomorphism (4) and the last is the well-known isomorphism. Let $u_1 : K_1(B) \rightarrow \text{Ext}_{\text{unital}}^{-1}(A, SB)$ be the composition

$$K_1(B) \longrightarrow K_1(Q(SB)) \xrightarrow{u} \text{Ext}_{\text{unital}}^{-1}(A, SB), \quad (11)$$

where the first map is the well-known isomorphism (the inverse of the one used in (10)) and second is the u -map as defined above, but with SB in place of B . Let $i_0^* : \text{Ext}^{-1}(A, SB) \rightarrow K_0(B)$ be the composition

$$\text{Ext}^{-1}(A, SB) \xrightarrow{i^*} \text{Ext}^{-1}(\mathbb{C}, SB) \longrightarrow K_0(B), \quad (12)$$

where the second map is the well-known isomorphism. Finally, let $u_0 : K_0(B) \rightarrow \text{Ext}_{\text{unital}}^{-1}(A, B)$ be the composition

$$K_0(B) \longrightarrow K_1(Q(B)) \xrightarrow{u} \text{Ext}_{\text{unital}}^{-1}(A, B), \quad (13)$$

where the first map is the well-known isomorphism. We have now all the ingredients to prove

Theorem 3. *The sequence*

$$\begin{array}{ccccc} K_0(B) & \xrightarrow{u_0} & \text{Ext}_{\text{unital}}^{-1}(A, B) & \longrightarrow & \text{Ext}^{-1}(A, B) \\ i_0^* \uparrow & & & & \downarrow i_1^* \\ \text{Ext}^{-1}(A, SB) & \longleftarrow & \text{Ext}_{\text{unital}}^{-1}(A, SB) & \xleftarrow{u_1} & K_1(B) \end{array}$$

is exact.

Proof. If we apply Lemma 2 with B replaced by SB we find that the sequence

$$\begin{array}{ccccc} \text{Ext}^{-1}(\mathbb{C}, SB) & & K_1(\alpha(A)' \cap Q(SB)) & & \\ i^* \uparrow & & \downarrow i^* & & \\ \text{Ext}^{-1}(A, SB) & \longleftarrow & \text{Ext}_{\text{unital}}^{-1}(A, SB) & \xleftarrow{u} & K_1(Q(SB)) \end{array} \quad (14)$$

is exact. Thanks to the commuting diagram (6) we can patch this sequence together with the sequence from Lemma 2 with the stated result. $\square$

REFERENCES

- [A] W. Arveson, *Notes on extensions of C^* -algebras*, Duke Math. J. **44** (1977), 329-355.
- [CH] J. Cuntz and N. Higson, *Kuiper's Theorem for Hilbert Modules*, Contemporary Mathematics **62**, 429-435, AMS, Providence, 1987.
- [M] J.A. Mingo, *On the contractibility of the unitary group of the Hilbert space over a C^* -algebra*, Integral Eq. Operator Theory **5** (1982), 888-891.
- [S] G. Skandalis, *Kasparov's bivariant K -theory and applications*, Exposition. Math. **9** (1991), no. 3, 193-250.
- [Th] K. Thomsen, *On absorbing extensions*, Proc. Amer. Math. Soc. **129** (2001), 1409-1417.

DEPARTMENT OF MECHANICS AND MATHEMATICS, MOSCOW STATE UNIVERSITY, LENINSKIE GORY,
MOSCOW, 119992, RUSSIA

E-mail address: `manuilov@mech.math.msu.su`

IMF, DEPARTMENT OF MATHEMATICS, NY MUNKEGADE, 8000 AARHUS C, DENMARK

E-mail address: `matkt@imf.au.dk`